Exhibit 1. Percent of Adults Under Age 65 with a Usual Source of Care, by State, 2012




Exhibit 2. Insured Adults More Likely to Have a Usual Source of Care, 2012

Percent of adults under age 65 with a usual source of care






Exhibit 3. Percent of Adults Under Age 65 with a Usual Source of Care, by Income and Insurance, 2012



Note: FPL = federal poverty level.

Exhibit 4. Percent of Older Adults Who Received Recommended Preventive Care, by State, 2012


Percent of adults ages 50 to 64



Note: Recommended care includes receipt of all of the following within a specific time frame given their age and sex: screenings for colon, breast, and cervical cancer, and flu shots.

Exhibit 5. Insured Older Adults More Likely to Receive Recommended Preventive Care, 2012


Percent of adults ages 50 to 64 receiving preventive care



Note: Recommended care includes receipt of all of the following within a specific time frame given their age and sex: screenings for colon, breast, and cervical cancer, and flu shots.


Exhibit 6. Percent of Older Adults Who Received Recommended Preventive Care, by Income and Insurance, 2012

Percent of adults ages 50 to 64



Notes: FPL = federal poverty level. Recommended care includes receipt of all of the following within a specific time frame given their age and sex: screenings for colon, breast, and cervical cancer, and flu shots.

Exhibit 7. Percent of Adults Under Age 65 Who Went Without Care Because of Cost, by Income and Insurance, 2012




Note: FPL = federal poverty level.

Exhibit 8. Percent of Older Adults Who Received Recommended Preventive Care, by Insurance, 2012



Insured Adults Ages 50 to 64



- 57%-68% (16 states + D.C.)
- 53%–56% (9 states)
- **50%–52% (11 states)**
- 44%–49% (14 states)

- 57%–69% (29 states + D.C.)
- 53%-56% (10 states)
- 50%-52% (8 states)
- 47%–48% (3 states)

Note: Recommended care includes receipt of all of the following within a specific time frame given their age and sex: screenings for colon, breast, and cervical cancer, and flu shots.