


# Exhibit 1. Continuously insured adults with private coverage or Medicaid rated the quality of their health care as excellent or very good at higher rates than did adults who were uninsured during the year.

Percent of adults ages 19–64



\* Difference is statistically significant from those with private coverage who were insured all year ( $p \leq 0.05$ ).


<sup>^</sup> Difference is statistically significant from those who were uninsured during the year ( $p \leq 0.05$ ). Percentages were adjusted for age, race, sex, health status, and income.

<sup>\*\*</sup> Excludes those who had not received health care in past 12 months.

Source: The Commonwealth Fund Biennial Health Insurance Survey, 2014.

## Exhibit 2. Adults with private coverage or with Medicaid who were insured all year reported that their doctor always or often knows their medical history at higher rates than did those who were uninsured during the year.

Percent of adults ages 19–64 who responded “always” or “often”



Adults ages 19–64 with a regular doctor or place of care

\* Difference is statistically significant from those with private coverage who were insured all year ( $p \leq 0.05$ ).

<sup>^</sup> Difference is statistically significant from those who were uninsured during the year ( $p \leq 0.05$ ). Percentages were adjusted for age, race, sex, health status, and income.


<sup>\*\*</sup> Excludes those who never tried to contact regular doctor by telephone.

<sup>\*\*\*</sup> Excludes those who never saw other doctors or went to another place for care.

Source: The Commonwealth Fund Biennial Health Insurance Survey, 2014.

# Exhibit 3. Adults with Medicaid coverage who were insured all year reported getting recommended preventive care services at higher rates than did those who were uninsured during the year.

Percent of adults ages 19-64



Note: Blood pressure checked in past two years (in past year if has hypertension or high blood pressure); cholesterol checked in past five years (in past year if has hypertension, heart disease, or high cholesterol); seasonal flu shot in past 12 months.


\* Difference is statistically significant from those with private coverage who were insured all year ( $p \leq 0.05$ ).

<sup>^</sup> Difference is statistically significant from those who were uninsured during the year ( $p \leq 0.05$ ). Percentages were adjusted for age, race, sex, health status, and income.

Source: The Commonwealth Fund Biennial Health Insurance Survey, 2014.

# Exhibit 4. Continuously insured adults with Medicaid coverage reported medical bill problems or having medical debt at lower rates than did those with private coverage and adults who were uninsured during the year.

Percent of adults ages 19–64




\* Difference is statistically significant from those with private coverage who were insured all year ( $p \leq 0.05$ ).

<sup>^</sup> Difference is statistically significant from those who were uninsured during the year ( $p \leq 0.05$ ). Percentages were adjusted for age, race, sex, health status, and income.

Source: The Commonwealth Fund Biennial Health Insurance Survey, 2014.

# Exhibit 5. Continuously insured adults with Medicaid coverage reported cost-related problems getting care at lower rates than did adults uninsured during the year.

Percent adults ages 19–64



\* Difference is statistically significant from those with private coverage who were insured all year ( $p \leq 0.05$ ).  
<sup>^</sup> Difference is statistically significant from those who were uninsured during the year ( $p \leq 0.05$ ). Percentages were adjusted for age, race, sex, health status, and income.